




Quality Measures

- Diabetes Care: LDL cholesterol monitoring testing
- Diabetes Care: Self-monitoring blood glucose testing
- Diabetes Care: Patients taking certain medications that had a test to monitor kidney function
- Diabetes Care: Patients that had at least 2 HbA1c blood sugar monitoring tests
- Diabetes Care: Patients that had a test to monitor kidney function
- Diabetes Care: Patients taking certain medications that had a serum potassium monitoring test
- Diabetes Care: Triglyceride monitoring testing
- Diabetes Care: Patients that had a HbA1c test in the last 12 months
- Diabetes Care: Patient(s) that did not have a diabetes related hospitalization in last 12 reported months
- Diabetes Care: Patient(s) that had at least one HbA1c test in the last 6 reported months
- Diabetes Care: Patient(s) that did not have a diabetes related emergency department encounter in last 12 reported months


Diabetes Care: LDL cholesterol monitoring testing

The percentage of adult diabetes patients with a LDL cholesterol in last 24 months.




Diabetes Care: Self-monitoring blood glucose testing

The percentage of adult diabetes patients taking insulin with evidence of self-monitoring blood glucose testing.



Diabetes Care: Patients taking certain medications that had a test to monitor kidney function

The percentage of diabetes patient taking a biguanide, ACE-inhibitor, or angiotensin II receptor antagonist that had a serum creatinine test to monitor kidney function.




Diabetes Care: Patients that had at least 2 HbA1c blood sugar monitoring tests

The percentage of diabetes patients that had at least 2 HbA1c blood sugar monitoring tests in last 12 reported months.




Diabetes Care: Patients that had a test to monitor kidney function

The percentage of adult diabetes patients that had serum creatinine test to monitor kidney function.




Diabetes Care: Patients taking certain medications that had a serum potassium monitoring test

The percentage of diabetes patients taking an ACE-inhibitor or angiotensin II receptor antagonist that had a serum potassium (K+) in last 12 reported months.




Diabetes Care: Triglyceride monitoring testing

The percentage of adult diabetes patients with a triglyceride test in last 12 months.




Diabetes Care: Patients that had a HbA1c test in the last 12 months

Adult diabetes patients ages 18-75 that had a HbA1c blood sugar test in the last 12 months.




Diabetes Care: Patient(s) that did not have a diabetes related hospitalization in last 12 reported months

Patient(s) that did not have a diabetes related hospitalization in last 12 reported months.



Diabetes Care: Patient(s) that had at least one HbA1c test in the last 6 reported months

Patient(s) that had at least one HbA1c test in the last 6 reported months.



Diabetes Care: Patient(s) that did not have a diabetes related emergency department encounter in last 12 reported months

Patient(s) that did not have a diabetes related emergency department encounter in last 12 reported months.

